



APS | BIOAG



Nutripore™

APPLICATION RATE AND TIMING:

Amenity/Grassland/Forage:

	Timing		Rate
Grass	Early Spring	Early spring at re-commencement of growth	20L/Ha
	Before onset of drought	Foliar application	20L/Ha
Split application recommended			
Amenity	Early Spring	Early spring at re-commencement of growth	20L/Ha
	Ongoing application	Reapply every 2 - 4 weeks thereafter	10L/Ha
Maize	4 - 6 leaf	For root development and good establishment	20L/Ha
	Prior to onset of mid season check	1st half of June when crop is at 6" to 12" (6L/Ha if first application missed)	20L/Ha
	At Plastic removal	Apply immediately after removing plastic	20L/Ha

Fruit & Vegetables:

Potatoes	Earlies & Bakers	4 days after 80% emergence	20L/Ha
		10 days after tuber initiation	20L/Ha
	Seed	4 days after 80% emergence	20L/Ha
Root Crops	Sugar Beet, Turnips, & Swedes	2 -4 pairs of leaves	20L/Ha
		At root swelling	20L/Ha
	Carrots	4 - 6 leaf stage	20L/Ha
		8 - 10 leaf stage	20L/Ha
		Root expansion stage	20L/Ha
Brassicas		4/5 leaf stage	20L/Ha
		Early development of button heads	20L/Ha
Soft Fruit	Strawberries	Early blossom	20L/Ha
	Raspberries	Fruit Set	20L/Ha
	Last Application	Immediately after harvest	20L/Ha
Top Fruit	From bud to fruiting	Apply regularly every 10 -14 days	5L/Ha
	Last Application	Immediately after harvest	5L/Ha



M +353 (0)87 828 8542
M +44 (0)739 351 2227
E leslie@apsbioag.ie

18 Dun na Rí, Carlow Road,
Athy, Co. Kildare, R14 FY05.

apsbioag.ie


Oifig Fiontair Áitiúil
Local Enterprise Office